AmeriChoice a UnitedHealth Group Company County of San Diego County Medical Services Program

CMS Medical Policies Issued July 2011



This document is periodically updated. Please visit the County Medical Services website for the most up-to-date issue at http://www.sdcounty.ca.gov/hhsa/programs/ssp/county_medical_services/in dex.html

For more information and/or comments regarding this document, please contact: Mayling Naputi, Medical Management Services Department Manager at (858) 495-1314 or email mayling_y_naputi@uhc.com

Table of Contents

ALLERGIC RHINITIS AND SINUSITIS	5
ARTHRITIS – OSTEOARTHRITIS	6
ASTHMA	7
BEE STING	8
BREAST CANCER - DIAGNOSIS AND TREATMENT	12
BREAST RECONSTRUCTION	17
BREAST REDUCTION SURGERY (REDUCTION MAMMOPLASTY)	18
BUNIONS	
BURNS – BURN CENTER POLICIES FOR MAJOR BURNS	22
CARDIAC NUCLEAR STRESS TESTING	23
CARDIAC STRESS TESTING (Medicated Non-Nuclear)	24
CARDIOLOGY	
CARPAL TUNNEL SYNDROME	
CHOLECYSTITIS AND CHOLELITHIASIS	27
CHRONIC FATIGUE SYNDROME	
COLONOSCOPY	
COLPOSCOPY - CERVICAL	
COMPRESSION STOCKINGS	32
CT OR MRI OF SPINE	
DENTAL	
DERMATOLOGY	38
DIABETIC SHOES	
DILATATION & CURETTAGE OF THE UTERUS (D&C)	
DME (DURABLE MEDICAL EQUIPMENT)	
DUPUYTREN'S CONTRACTURE	
ESOPHOGASTRODUODENOSCOPY (EGD)	
ENDOMETRIAL ABLATION	
EPIDURAL STEROID INJECTION	
EPILEPSY (SEIZURE DISORDER) VAGUS NERVE STIMULATION (VNS)	
ERYTHROPOIETIN (RECOMBINANT GROWTH FACTOR), EPOGEN, PROCRIT.	
	52
GENETIC COUNSELING	
GYNECOMASTIA	
HEADACHE - MIGRAINES	
HEADACHE	
HEARING LOSS	
HEMORRHOIDECTOMY	
HEPATITIS B & C	
HERNIAS – SURGICAL CORRECTION	
HIRSUTISM	61

HORMONE THERAPY	62
HYDROCELE	
HYPERPARATHYROIDISM (CAUSING HYPOCALCAEMIA)	64
HYPOGONADISM AND TESTOSTERONE REPLACEMENT	
HYSTERECTOMY	
HYSTERECTOMY WORKSHEET	
HYSTEROSCOPY	69
INCONTINENCE	
INFLUENZA (INCLUDING SWINE FLU)	71
INSULIN PUMP	
LAMINECTOMY (SPINE SURGERY)	73
LAPAROSCOPY (GYNECOLOGY)	
MAMMOGRAPHY	
MEDICAL TRANSPORTATION	76
MRI	78
MRI OF KNEE	79
NEPHROLOGY	81
OCULAR DISEASE - REFERRAL FOR VISION LOSS OR EYE PAIN	82
OPHTHALMOLOGY	
OPHTHALMOLOGY - CATARACTS	85
OPHTHALMOLOGY - GLAUCOMA	86
OPHTHALMOLOGY - PTERYGIUM	87
OPHTHALMOLOGY - FOLLOW-UP EXAMINATION GUIDELINES	88
OPTOMETRY- COVERAGE FOR GLASSES	90
ORGAN TRANSPLANTATION	91
OTITIS MEDIA	92
PAIN MANAGEMENT	93
PEPTIC ULCER DISEASE	
PET SCAN (POSITRON – EMISSION TOMOGRAPHY)	95
PHYSICAL THERAPY, OCCUPATIONAL THERAPY, SPEECH THERAPY	96
PLANTAR FASCIITIS	97
PODIATRY	
PROSTATECTOMY	
RHEUMATOLOGY - INCLUDING RHEUMATOID ARTHRITIS	
SECOND OPINION	
SEPTOPLASTY	
SINUSITIS – ACUTE AND CHRONIC	
SLEEP APNEA	104
THYROID DISEASE	
THYROIDECTOMY	
TEMPOROMANDIBULAR JOINT DISORDER-TMJ	
TONSILLECTOMY AND ADENOIDECTOMY (T & A)	110

TRIGGER FINGER	111
TYMPANOPLASTY	112
TYMPANOTOMY	113
VARICOSE VEINS	
VERTIGO	115
WORK HISTORY INFORMATION	117
WOUND MANAGEMENT	118

ALLERGIC RHINITIS AND SINUSITIS

The CMS program does not cover treatment or referral for common allergic rhinitis. Treatment or referral is covered only for disease which interferes with the ability to function and work.

Criteria for Authorization

Patient History (two of three)

- chronic symptoms, at least 3 days per week
- facial pain
- chronic purulent discharge

Physical Exam (two of three)

- facial tenderness
- green/yellow discharge
- swelling and polypoid changes in the nose

Medication failure (all three)

- decongestants and/or antihistamines
- antibiotics for 6 weeks
- nasal steroids and/or nasal Cromolyn Sodium

X-Rays

 sinus imaging (plain films or CT scan) showing evidence of infection

ARTHRITIS - OSTEOARTHRITIS

Criteria for Authorization

Referral to Orthopedic Surgery is only allowable if the patient requires surgery to function at work or with daily activities. Joint replacement for Osteoarthritis is not a CMS benefit. Physical therapy is approved only if rehabilitation is necessary. Joint replacement is not a covered benefit [special circumstances may be considered for approval replacement, considering work history, age of patient, etc.]

Patient History (three of four)

- restriction of daily activities
- interferes with current work
- failure to respond to medications 3 month trial
- failure to respond to physical therapy (Orthopedic referral)

AND

Physical Exam (two of four)

- tenderness with movement
- decrease range of motion
- muscle wasting
- deformity

AND

Imaging

evidence of moderate to severe joint changes

ASTHMA

Mild intermittent, mild persistent and moderate persistent asthma are managed at the primary care level. Severe asthma, defined as requiring continuous systemic steroid therapy and a history of hospitalization, should be referred to an allergy or pulmonary specialist. Desensitization is covered by CMS only for asthma which interferes with function or work.

Criteria for Authorization

Patient History (one of three)

- life threatening
- asthma not responding to maximum medical therapy
- multiple ER visits, > 2 per year, or hospitalization

Treatment failure (two of four)

- b-agonists, including long acting
- theophylline
- cromolyn sodium
- inhalation corticosteroids for 3 or more months

Tests

pulmonary function testing which shows severe reversible disease

BEE STING

The CMS program covers Bee Sting Allergy kits for a history of definite systemic allergic reaction to bee stings. Referral for consultation and desensitization is based on the following criteria.

Criteria for Authorization

Patient History (three)

- respiratory distress, acute urticaria or hypotension after a bee sting (history of anaphylaxis)
- reaction of bee sting is remote from the local reaction, at least 6 inches from sting
- personal risk at work or at home for bee sting exposure

Physical Exam (not required if history is clear or reaction documented by past medical records).

Evidence of allergic reaction remote from the site of the sting, including hives (urticaria), respiratory distress or hypotension.

BONE GROWTH STIMULATOR

Although bone stimulators have been used in a variety of clinical settings, the following requirements, adapted from the MediCal program (April 2008 criteria), limit the utilization of osteogenesis stimulator devices.

Criteria for Authorization

Spinal Electrical Device (cervical, thoracic, lumbar, and sacral vertebrae)

(one of the following must be met)

- Failed spinal fusion and a minimum of nine months have elapsed since the last surgery, or
- Following a multi-level spinal fusion surgery involving three or more vertebrae (for example, L3-5, L4-S1, etc.), or
- Following spinal fusion surgery where there is a history of a previously failed spinal fusion at the same site, or
- The patient has one or more risk factors for high risk of spinal fusion failure such as smoking, obesity (BMI>35), diabetes, renal disease, alcoholism, grade II or worse Spondylolisthesis, or other metabolic disease where bone healing is poor

Note: The device should be applied within 30 days as an adjunct to spinal fusion surgery. The patient should use the device for at least two hours per day, and the treatment period continued for nine months (270 consecutive days).

Low-Intensity Ultrasound Device

 Nonunion of a fracture other than the skull or vertebrae, documented by a minimum of 2 sets of radiographs, each including multiple views of the fracture site, separated by a minimum of 90 days, obtained prior to starting treatment with the osteogenesis stimulator, with a written

BONE GROWTH STIMULATOR (Continued)

interpretation by a physician stating that there has been no clinically significant evidence of fracture healing between the two sets of radiographs, AND the fracture is not tumor-related.

- OR one of the following for new fractures (not tumor-related):
 - Fresh (<7 days), closed or grade I open, tibial diaphyseal fractures, or
 - Fresh (<7 days), closed fractures of the distal radius (Colles fracture).

Note: An ultrasonic osteogenesis stimulator may not be used concurrently with other noninvasive stimulators.

- If surgery has been performed in an attempt to induce healing of the bone fracture or nonunion via bone graft or internal fixation, the six or twelve month duration begins from the date of the surgery. The surgery date and procedure performed should be included with the TAR.
- Although the pulse generator is returned to the manufacturer, the device is billed as a purchase because the manufacturer does not provide the generator separately from the coils. Whenever possible, attempt to rent the device for one fee which will cover any periods up to one year.
- If the device is approved as a rental, authorize for a one year period. Any reauthorization request should be denied, because if the OBS has failed to produce complete union within one year's time, the device is not effective in that case.

BONE MINERAL DENSITY (BMD) TESTING

The CMS program covers only diagnostic evaluation to confirm the presence of suspected disease and provide critical treatment. Screening BMD is not covered. There must be evidence of likely or present osteoporosis or other metabolic bone disease.

<u>Criteria for Authorization</u>

- Patient has vertebral abnormalities as demonstrated by X-ray to be indicative of osteoporosis, low bone mass (osteopenia), or vertebral fracture.
- Glucocorticoid therapy equivalent of 7.5 mg of prednisone or greater per day for 3 months or longer, or the equivalent of 5 mg of prednisone or greater for 6 months or longer.
- Patient has hyperparathyroidism.
- Patient is being monitored to assess the response to or efficacy of an FDA approved osteoporosis drug therapy.

BREAST CANCER - DIAGNOSIS AND TREATMENT

The CMS program follows California law for the diagnosis and treatment of breast cancer.

The people of the State of California do enact as follows:

- SEC. 1. Section 1367.6 of the Health and Safety Code is repealed.
- SEC. 2. Section 1367.6 is added to the Health and Safety Code, to read:
- 1367.6. (a) Every health care service plan contract, except a specialized health care service plan contract, that is issued, amended, delivered, or renewed on or after January 1, 2000, shall provide coverage for screening for, diagnosis of, and treatment for, breast cancer.
- (b) No health care service plan contract shall deny enrollment or coverage to an individual solely due to a family history of breast cancer, or who has had one or more diagnostic procedures for breast disease but has not developed or been diagnosed with breast cancer.
- (c) Every health care service plan contract shall cover screening and diagnosis of breast cancer, consistent with generally accepted medical practice and scientific evidence, upon the referral of the enrollee's participating physician.
- (d) Treatment for breast cancer under this section shall include coverage for prosthetic devices or reconstructive surgery to restore and achieve symmetry for the patient incident to a mastectomy. Coverage for prosthetic devices and reconstructive surgery shall be subject to the co-payment, or deductible and coinsurance conditions, that are applicable to the mastectomy and all other terms and conditions applicable to other benefits.

BREAST CANCER - DIAGNOSIS AND TREATMENT (Continued)

- (e) As used in this section, "mastectomy" means the removal of all or part of the breast for medically necessary reasons, as determined by a licensed physician and surgeon.
- (f) As used in the section, "prosthetic devices" means the provision of initial and subsequent devices pursuant to an order of the patient's physician and surgeon.
- SEC. 3. Section 1367.65 of the Health and Safety Code is amended read:
- 1367.65 (a) On or after January 1, 2000, every health care service plan contract, except a specialized health care service plan contract, that is issued, amended, delivered, or renewed shall be deemed to provide coverage for mammography for screening or diagnostic purposes upon referral by a participating nurse practitioner, participating certified nurse midwife, or participating physician, providing care to the patient and operating within the scope of practice provided under existing law.
- (b) Nothing in this section shall be construed to prevent application of co-payment or deductible provisions in a plan, nor shall this section be construed to require that a plan be extended to cover any other procedures under an individual or a group health care service plan contract. Nothing in this section shall be construed to authorize a plan enrollee to receive the services required to be covered by this section if those services are furnished by a nonparticipating provider, unless the plan enrollee is referred to that provider by a participating physician, nurse practitioner, or certified nurse midwife providing care.
- SEC. 4. Section 10123.8 of the Insurance Code is repealed.
- SEC. 5. Section 10123.8 is added to the Insurance Code, to read:
- 10123.8 (a) Every policy of disability insurance that provides coverage for hospital, medical, or surgical expenses, that is issued, amended, delivered, or renewed on or after January 1, 2000, shall provide coverage for screening for, diagnosis of, and treatment for, breast cancer.

BREAST CANCER - DIAGNOSIS AND TREATMENT (Continued)

- (b) No policy of disability insurance that provides coverage for hospital, medical, or surgical expenses shall deny enrollment or coverage to an individual solely due to a family history of breast cancer, or who has had one or more diagnostic procedures for breast disease but has not developed or been diagnosed with breast cancer.
- (c) Every policy of disability insurance shall cover screening and diagnosis of breast cancer, consistent with generally accepted medical practice and scientific evidence, upon the referral of the insured's participation physician.
- (d) Treatment for breast cancer under this section shall include coverage for prosthetic devices or reconstructive surgery to restore and achieve symmetry for the patient incident to a mastectomy. Coverage for prosthetic devices and reconstructive surgery shall be subject to the deductible and coinsurance conditions applied to the mastectomy and all other terms and conditions applicable to other benefits.
- (e) As used in this section, "mastectomy" means the removal of all or part of the breast for medically necessary reasons, as determined by a licensed physician and surgeon.

As used in this section, "prosthetic devices" means the provision of initial and subsequent devices pursuant to an order of the patient's physician and surgeon.

(f) For the purposes of this section, disability insurance does not include accident only, credit, disability income, specified disease and hospital confinement indemnity, coverage of Medicare services pursuant to contracts with the United States government, Medicare supplement, long-term care insurance, dental, vision, coverage issued as a supplement to liability insurance, insurance arising out of a workers' compensation or similar law, automobile medical payment insurance, or insurance under which benefits are payable with or without regard to fault and that is statutorily required to be contained in any liability insurance policy or equivalent self-insurance.

BREAST CANCER - DIAGNOSIS AND TREATMENT (Continued)

SEC. 6. Section 10123.81 of the Insurance Code is amended to read:

10123.81 On or after January 1, 2000, every individual or group policy of disability insurance of self-insured employee welfare benefit plan that is issued, amended, or renewed, shall be deemed to provide coverage for at least the following, upon the referral of a nurse practitioner, certified nurse midwife, or physician, providing care to the patient and operating within the scope of practice provided under existing law for breast cancer screening or diagnostic purposes:

- a) A baseline mammogram for women age 35 to 39, inclusive.
- b) A mammogram for women age 40 to 49, inclusive, every two years or more frequently based on the women's physician's recommendation.
- c) A mammogram every year for women age 50 and over.

Nothing in this section shall be construed to require an individual or group policy to cover the surgical procedure known as mastectomy or to prevent application of deductible or copayment provisions contained in the policy or plan, nor shall this section be construed to require that coverage under an individual or group policy be extended to any other procedures.

Nothing in this section shall be construed to authorize an insured or plan member to receive the coverage required by this section if that coverage is furnished by a nonparticipating provider, unless the insured or plan member is referred to that provider by a participating physician, nurse practitioner, or certified nurse midwife providing care.

SEC. 7. No reimbursement is required by this act pursuant to Section 6 of Article XIII B of the California Constitution because the only costs that may be incurred by a local agency or school district will be incurred by a local agency or school district will be incurred because this act creates a new crime or infraction, eliminates a crime or infraction, or changes the penalty for a BREAST CANCER – DIAGNOSIS AND TREATMENT (Continued)

BREAST CANCER - DIAGNOSIS AND TREATMENT (Continued)

crime or infraction, within the meaning of Section 17556 of the Government Code, or changes the definition of a crime within the meaning of Section 6 of Article XIII B of the California Constitution.

BREAST RECONSTRUCTION

The CMS program covers breast reconstruction only in relation to breast cancer treatment and following or concordant with breast cancer surgery. In rare instances, breast reconstruction may be approved for removal of prosthesis if it is extruding and interferes with the ability to work.

Criteria for Authorization

- following or concordant with breast cancer surgery
- treatment associated abnormalities or deformities
- removal of prosthesis for fibrosis or extrusion (work history required)
- replacement of prosthesis if mastectomy due to breast cancer
- nipple reconstruction (for breast cancer)
- breast prosthesis and bras (2), with replacements covered every two years

BREAST REDUCTION SURGERY (REDUCTION MAMMOPLASTY)

The CMS program covers breast reduction surgery only if it is designated medically necessary, in relation to the therapeutic treatment of a medical condition. Reduction mammoplasty is the removal of breast tissue to reduce size and weight of mammary tissue.

Breast reduction surgery is considered not medically necessary for the following conditions: poor posture, breast asymmetry, pendulousness, problems with clothes fitting properly and nipple-areola distortion and/or psychological considerations.

Mastoplexy or breast lift is a cosmetic reshaping of the breast by removal of skin with or without a small amount of breast tissue and is never covered by CMS.

Criteria for Authorization

Criteria for Authorization for a Consultation for Therapeutic Reduction Mammoplasty

- Evidence of breasts large enough to cause pain or intertriginous dermatitis
 - o Bra size D cup or larger
 - Shoulder bra strap discomfort and demonstrable severe shoulder grooves and/or intractable dermatitis due to bra strap pressure
- AND evaluation of upper back (thoracic and cervical) severe chronic pain (1 year or greater duration) including:
 - Diagnostic testing to evaluate the causes of pain.
 - Evaluations by appropriate consultant(s) from the specialty area(s) of orthopedics, neurology, rheumatology, and/or pain management if the member's PCP requests further assessment of the cause of pain.
- AND documentation of at least 3 months of a reasonable trial of conservative therapy including all of the following:

BREAST REDUCTION SURGERY (REDUCTION MAMMOPLASTY) (Continued)

- A reasonable trial of NSAIDs (nonsteroidal anti-inflammatory drugs) pain medications and/or muscle relaxants without relief of symptoms.
- o Physical therapy, exercise program, and the use of properly fitting undergarments.
- o BMI less than 30.
- o There is a reasonable prognosis of symptom relief with reduction mammoplasty.

Criteria for Authorization for Therapeutic Reduction Mammoplasty

- Documentation of all of the following is required to substantiate medical necessity for therapeutic reduction mammoplasty:
- A significant Therapeutic Tissue Reduction/Ratio
 - o The appropriate amounts (in grams) of breast tissue must be anticipated for removal from each breast, which is based on the patient's total body surface area (BSA) in meters squared. See Table for BSA values to the minimum amount (weight) of breast tissue to be removed per breast.
- AND excessively large pendulous breasts out of proportion to the rest of the individual's normal body structure as demonstrated by measurement, e.g., a suprasternal notch to nipple measurement of greater than or equal to 27cm (average range is 20-24cm)

BREAST REDUCTION SURGERY (REDUCTION MAMMOPLASTY) (Continued)

Minimum Weight of Breast Tissue Removed, per Breast, as a Function of Body Surface Area Schnur Sliding Scale

Body Surface Area (meters squared)	Minimum weight of tissue to be removed per breast (grams)
1.35	199
1.40	218
1.45	238
1.50	260
1.55	284
1.60	310
1.65	338
1.70	370
1.75	404
1.80	441
1.85	482
1.90	527
1.95	575
2.00	628
2.05	687
2.10	750
2.15	819
2.20	895
2.25	978
2.30 or greater	>= 1000

Calculation: BSA=square root of {(height in inches x weight in pounds) / 3131}

BUNIONS

Criteria for Authorization

Patient History

• symptomatic bunions which interfere with daily function or work.

AND

Physical Exam

marked deformity is present

AND

Radiology

- x-ray confirmation of severe deformity
- hallux valgus angle greater than 40 degrees
- marked lateral deviation
- some overlapping of second toe
- subluxation of lateral sesamoid bone

AND

Work History

BURNS - BURN CENTER POLICIES FOR MAJOR BURNS

The CMS Program covers up to 5 days of UCSD Burn Center level of treatment and reimbursement. Any extension of this coverage requires Medical Director review and approval.

Criteria for Authorization

- patient has 30% or more 3rd degree burns, or
- patient has inhalation burns requiring intubation, or
- level of care required is intensive for both medical and burn care

Rule of Nines

Wallace's Rule of Nines provides an approximation of the area of skin burned. It divides the body into units of surface are divisible by nine, with the exception of the perineum. Charts are available in most burn units.

The following are the respective percentages of an adult body surface area:

- head and neck front and back 9%
- each upper limb total front and back 9%
- chest and Abdomen front 18%
- back side, thorax and low back 18%
- perineum 1 %
- each lower limb total for front and back 18%

CARDIAC NUCLEAR STRESS TESTING

Criteria for Authorization

A cardiac thallium scan is done to evaluate the chambers and structure of the heart while exercising. It is used to evaluate circulation to the heart muscle. A cardiac thallium is more accurate in evaluation evidence of coronary heart disease than a regular exercise stress test.

Patient History Required:

- prior treadmill testing
- abnormal EKG 1mm or greater deviation from normal baseline
- bundle branch block
- unstable angina
- female > 45 or surgical menopause with chest pain or anginal equivalent

CARDIAC STRESS TESTING (Medicated Non-Nuclear)

Exercise Cardiac Stress Testing is done on all patients except those unable to exercise on the treadmill. Referral for Cardiac Stress Testing is based on a clear risk for coronary artery disease. Screening Cardiac Stress Testing is not approved by the CMS program. The patient must have two or more cardiac risk factors (age over 40, positive family history, smoking, hypertension, hyperlipidemia, diabetes, and obesity) to qualify for medicated non-nuclear cardiac stress testing.

Medicated Cardiac Stress Testing (Adenosine, Dipyridamole) is approved based on the following:

<u>Criteria for Authorization</u>

History (one of three)

- incapable of exercising on a treadmill
 - difficulty maintaining balance
 - excessive obesity
- reactive airway disease risk of bronchospasm
- frail and elderly

AND

Conjunction Testing

- echocardiography
- nuclear scan

CARDIOLOGY

Referral for Cardiology is for critical need only, not for screening. Patients with active heart disease, such as coronary artery disease, heart failure, cardiomyopathy or cardiac arrhythmia may be followed by a cardiologist. Cardiac consultation is approved with clear evidence of a cardiac condition, and for an acutely ill or unstable patient. The initial evaluation for cardiac disease is done by the primary care physician (PCP). The PCP may order without a TAR, on a supplemental form, the following studies: EKG, Exercise EKG (stress test), Echocardiogram, Stress Echocardiogram and Holter monitor.

Criteria for Authorization

History (two of four)

- chest pain on exertion
- shortness of breath on exertion
- major risk factors for heart disease, including diabetes mellitus, hypertension, dyslipidemia, smoking, family history, obesity, age over 40 (male) and over 50 (female)
- new onset of weakness and fatigue

Physical Exam (one of four)

- cardiac murmur
- abnormal heart sounds
- peripheral edema
- jugular venous distention

AND

Testing (one of five)

- abnormal EKG
- abnormal echocardiography
- abnormal exercise EKG
- cardiomegaly by chest x-ray or echocardiogram
- abnormal holter monitor

CARPAL TUNNEL SYNDROME

Most patients with carpal tunnel syndrome improve and recover in the primary care setting. For many patients, carpal tunnel syndrome is an overuse injury and a change in work position and rest resolve the problem. For patients with persistent carpal tunnel syndrome despite rest and wrist splinting, referral to an orthopedic surgeon for injection or surgery may be necessary. Referral to Neurology for nerve conduction study is unnecessary unless the diagnosis is uncertain.

Criteria for Authorization

Patient History (one of two required)

- failed three months of conservative management
 - regular use of NSAIDs and night splints
 - trial of changed work positioning
- interferes with A.D.L

AND

Physical Exam (one of three required)

- positive tinel's and/or phalen's test
- atrophy or weakness of the thenar muscles
- documented nerve impairment on nerve conduction velocity testing (done only if the physical exam is uncertain)

AND

Work History (a procedure will help the patient continue or return to work)

CHOLECYSTITIS AND CHOLELITHIASIS

Criteria for Authorization

The CMS program covers cholecystectomy only for the removable of symptomatic gallstones causing clinical obstruction or infection. The CMS program does not cover surgery for asymptomatic gallstones

History (one of four)

- the diagnostic tests verify the presence of gallstones
- history of jaundice
- two documented episodes of abdominal colic or RUQ pain
- the presence of nausea/vomiting, chills and fever, leukocytosis

AND

Physical (one of two)

- the patient has abdominal guarding/tenderness
- a mass in the RUQ

AND

Diagnostic Tests

ultrasound or CT scan documents presence of gallstones

CHRONIC FATIGUE SYNDROME

Criteria for Authorization

History (four of four)

- severe unexplained fatigue for > 6 months
- functionally impaired
- identifiable date of onset
- unrelated to psychological stress

AND

Symptoms (three of eight)

- memory or concentration complaints
- sore throat
- tender lymph nodes
- muscle pain
- multi-joint pain
- new pattern of headaches
- unrefreshing sleep
- postexertional malaise lasting more than 24 hours

AND

Treatment (five of five)

- judicious use of medication to ameliorate symptoms
- graded exercise or rehabilitation measures
- hypothyroidism has been ruled out
- depression has been ruled out or treated
- family history of colon cancer
 - three 1° relatives q 3-5 years from age 20
 - one or two 1° relatives q 3-5 years from age 40

COLONOSCOPY

Criteria for Authorization

CMS does not cover screening colonoscopy. Any one of the following indications must be met.

History (at least one present)

- positive stool culture or O&P study
- unexplained iron deficiency anemia
- acute diarrhea following recent antibiotic therapy
- melena normal UGI endoscopy
- rectal bleeding unexplained
- abnormal x-ray findings, mass, lesion or ulceration
- chronic diarrhea
- ulcerative colitis
- crohn's disease

OR

Therapeutic (at least one present)

- excision of polyps
- removal of foreign body
- dilatation of stricture
- control active bleeding

OR

Surveillance (at least one present)

- with colon polyps every 3-5 years, if large (greater than 2 cm), may repeat in 3-6 months, if multiple adenomas, repeat at 1 and 4 years.
- following polypectomy 1 year, then 1-3 year intervals
- following removal of colon cancer, 6 months, 1 year, q 2-3 years
- ulcerative colitis q 1-2 years after 8th year when stable

COLONOSCOPY (Continued)

- left sided colitis q 1-2 years after 15th year when stable
- family history of colon cancer
 - three 1° relatives q 3-5 years from age 20
 - one or two 1° relatives q 3-5 years from age 40

COLPOSCOPY - CERVICAL

Criteria for Authorization

Colposcopy is performed to evaluate abnormal Pap Smears and to allow for guided cervical biopsies. Not all atypical Pap smears require Colposcopy, and Colposcopy should not be routinely repeated if the cervical abnormalities are minor. Repeat Pap smears are an acceptable way to monitor mild cervical pathology, especially if HPV testing is negative.

Colposcopy is indicated for cervical cytology demonstrating:

- ASCUS (Atypical Squamous Cells of Undetermined Significance)
 - 2 or more Pap reports are abnormal
 - Positive HPV testing
 - Suspicious cervical lesion
 - ASCUS persists in repeat pap 3-6 months after infections are treated
- SIL (Moderate Dysplasia, CIN I)
 - visible abnormality
- HSIL (Moderate and Severe Dysplasia, CIN II, CIN III)
- Other suspicious lesion

COMPRESSION STOCKINGS

<u>Criteria for Authorization</u>

Compression stockings are used for a variety of conditions: dependent edema, chronic venous insufficiency, recurrent leg ulcers and for wound management. CMS will approve compression stockings only when critically necessary to restore or maintain function in the patient, such as allowing the patient to work. When approved, two pair of stockings are allowable with a renewal no sooner than six months.

CT OR MRI OF SPINE

Criteria for Authorization

Advanced imaging of the spine should be performed for specific indications and not simply because of pain. In the presence of chronic pain, advanced imaging is done only if there are symptoms of neurologic impairment or suspicion of a lesion in the bone. CT is done to evaluate the bone tissue, and MRI is preferred for looking at the spinal cord and nerves. Indications include:

- suspected fractures and dislocations (not clear by plain x-rays) (CT)
- disk herniations causing neurologic signs or symptoms (MRI)
- previously documented spinal stenosis (MRI)
- previous spinal surgery and demonstration of nonunion on x-ray or positive Bone Scan (CT)
- significant trial of conservative therapy including anti-inflammatory medications and physician supervised home exercise/physical therapy (MRI)
- chronic pain in a patient at risk for cancer (CT)
- localized tenderness of a vertebral body suggesting osteomyelitis(CT)

DENTAL

Criteria for Authorization

The County Medical Services Dental Program is designed to provide EMERGENCY dental treatment to alleviate a patient's IMMEDIATE source of dental pain. Dental clinics will provide emergency dental care based on the CMS basic dental service list; all other procedures/treatments must be medically indicated and require prior approval.

Emergency Care Only (for acute pain)

- site specific x-rays
- urgent extractions
- palliative Restoration fillings
- treatment of existing dental disease that has a significant effect on the patient's chronic disease, nutritional or employment status

Endodontics (Root canal treatment): TAR Required

- Anterior teeth (6-11 and 22-27) are covered. Other teeth may be approved under circumstances of persistent symptoms and need to maintain nutrition. The final composite restoration is covered under the basic encounter fee. The composite restoration only is covered (Patient may pay for a crown).
- Necessary to ensure adequate medical treatment (documentation of health condition required).

Not Covered:

- routine dental examinations, x-rays, cleaning, or prophylaxis
- restoration of asymptomatic teeth
- comprehensive periodontal treatment
- permanent crowns and bridges
- dental prosthesis not meeting the above criteria
- orthodontia

DENTAL (Continued)

- elective services and/or medications not required to treat a potentially disabling or life-threatening illness or condition
- immediate dentures and immediate partial dentures

The Following should be submitted with Dental TARS:

illustration of mouth with teeth numbered

<u>Specific Dental Prostheses:</u>

Partial and Full Dentures are indicated for critical function purposes and not for cosmetics. In most circumstances, only the anterior teeth (partial denture) are covered.

Partial Dentures are indicated:

- to replace recently extracted teeth only after adequate healing to allow pursuit of employment (documentation required)
- evaluation of the remaining teeth (must be good dentition, with enough posterior teeth to hold the partial denture
- anterior teeth only (6-11, 22-27)
- repair or replacement of an existing appliance one time in 12 months (once a year) is covered
- completion of Work history form is required

Full dentures are indicated for:

- an edentulous patient, to replace extracted or missing teeth
- needed to maintain nutritional status, maintain function (including speech), and secure employment
- repair or replacement of an existing appliance one time in 12 months (once a year)
- completion of Work history form is required

Aveleoplasty:

- if patient does not already have dentures, need a work history.
- approve for re-fit when patient has existing dentures

DENTAL (Continued)

Stainless steel:

• crowns are only covered if a posterior tooth has no integrity, but the remaining teeth are in good shape.

Oral Surgery: CMS only covers Oral Surgery for critically necessary services (TAR required):

- wisdom teeth (#1, 16, 17, 32) only symptomatic 3rd molars are covered. Soft tissue impaction can be extracted in the dental clinic. Authorization is not needed.
- oral lesions to remove or biopsy for cancer

San Diego County CMS Dental Program Work History Information

The CMS Program policy limits dental services, specifically stay-plates and dentures. We require specific information from the patient to determine if the requested service meets all of the CMS Program guidelines to be a covered service. Your cooperation is appreciated. **ALL questions must be answered and the form attached to the request for dental replacements.**

Date Sent:					
Patient Name:	SSN:	7			
Phone Number:	DOB:				
What kind of dental service do you need?	-				
2. When were your teeth extracted? Month	Year				
3. What kind of work do you do when you are work	king?				
4. Are you currently employed?	[]Yes []No				
5. Are you currently Receiving State Disability?	[]Yes []No				
6. Are you currently receiving workers compensation	ion? []Yes []No				
7. Date you last worked?					
IF YOU ARE CURRENTLY UNEMPLOYED:					
Why did you leave your last job?					
2. Have you applied for or been offered employment in the past (6) months? []Yes [] N					
3. Have you recently been turned down for a job because of this medical condition? []Yes [] N					
TELL US WHO YOUR CURRENT EMPLOYER IS OFFERED YOU EMPLOYMENT	OR ABOUT THE COMPANY WHO HAS				
Name of Company:					
Person to Contact: Phone:					
If you are currently employed you can speed up the employer and send a letter on <u>business letterhead.</u> employment and how this condition affects your abin history and send them to: CMS Pro ATTN: Authorizati	. This letter should tell us about your bility to do your job. Attach the letter to this work rogram tion Coordinators				
PO Box 9 San Diego, 0 I authorize the CMS Program to contact the person	CA 92193				
information presented.					
Patient Signature:	Date:				
CA4CA4 I' I D - I' - i	07 - [110				

DERMATOLOGY

Criteria for Authorization

Referral and Office biopsy is allowed for the following:

- possible Melanoma
- basal Cell Cancer (BCC)
- squamous Cell Cancer
- other Malignant neoplasm
- dysplastic nevi

Referral for Certain Conditions:

Pruritis: (Itching): Most itching in primary care is due either to excessive use of soap (dry skin) or neurodermatitis. The primary care physician is able to evaluate and treat most causes of pruritis, including primary and secondary conditions. Referral is approved to Dermatology only after a thorough primary care evaluation is done and the cause is uncertain or the treatment is ineffective. The patient history, physical findings and previous treatments must be well documented.

Rash: Most skin rashes are diagnosed and treated in primary care. Referral to Dermatology is approved when the diagnosis and treatment remain uncertain. The patient history, physical findings and previous treatments must be well documented.

Acne: The CMS program does not cover the treatment of mild or moderate acne. Referral to Dermatology would only be allowed for severe, cystic, inflammatory acne. The patient history, physical findings and previous treatments must be well documented.

Psoriasis: Referral to Dermatology is approved only for psoriasis in multiple areas which is actively inflammatory and unstable. The patient history, physical findings and previous treatments must be well documented.

Treatments for psoriasis which are not on formulary,

DERMATOLOGY (Continued)

or procedures such as PUVA, require prior authorization.

Actinic Keratosis: Most actinic keratosis is managed by primary care.

Referral to Dermatology is allowed for extensive disease on exposed areas. The patient history, physical findings and previous treatments must be well documented. Treatments which are not on formulary require prior

authorization.

DIABETIC SHOES

Diabetic shoes are frequently recommended to protect the feet of patients with diabetic neuropathy. The CMS program does not cover customized diabetic shoes for patients with diabetes or any other medical condition unless a specialist is able to indicate that such treatment is critically necessary for work or life function. Indications for diabetic shoes include patients with an amputation, with a chronic wound or ulcer, or in someone with an extensive history of such.

DILATATION & CURETTAGE OF THE UTERUS (D&C)

Criteria for Authorization

D & C is performed less commonly today with better procedures for evaluating the tissue in the uterus. For diagnostic purposes, endometrial biopsy, hysteroscopy and ultrasound are often used when D & C had been done in the past.

Patient History (either one of these present)

- excessive bleeding with a suspicion of tissue present
- post-menopausal bleeding to evaluate for endometrial cancer
- recurrent post-menopausal bleeding (for treatment, other tests are performed for diagnosis)

Addendum:

- there is no indication for performing a D&C in an adolescent
- heavy bleeding is usually better treated with hormones
- office endometrial biopsy is the first step to evaluate for endometrial cancer, and hysteroscopy is preferable to a D & C as the second test
- bleeding associated with pregnancy is not a covered CMS benefit

DME (DURABLE MEDICAL EQUIPMENT)

Durable medical equipment is reusable medical equipment such as walkers or wheelchairs. CMS covers durable medical equipment which is medically necessary when prescribed by a doctor or treating practitioner to be used in the patient's home.

Specifically, DME is defined as equipment that:

- can withstand repeated use;
- is used to serve a medical purpose
- is not useful to an individual in the absence of illness, injury, functional impairment, or congenital anomaly; and
- is appropriate for use in or out of the patient's home.
- not considered disposable, with the exception of ostomy bags
- is necessary to preserve bodily functions essential to activities of daily living; and
- provides therapeutic benefits to a patient in need because of certain medical conditions and/or illnesses.

All requirements of the definition must be met before an item can be considered to be durable medical equipment.

<u>Criteria for Authorization</u>

Medically necessary DME is covered only when:

- medically necessary for the specific member's medical condition or functional limitation
- equipment does not have significant non-medical uses
- is not duplicative of the function of another piece of equipment or device already provided for the member
- is intended for exclusive use of a CMS member
- is ordered and/or prescribed by a CMS provider practicing within their scope of practice
- is lowest cost DME item necessary to meet patient's needs

DME (DURABLE MEDICAL EQUIPMENT) (Continued)

DME normally does not include:

- disposable medical supplies
- devices or equipment used for environmental control (e.g., electric air cleaners, room heaters) or to enhance the environmental setting (e.g., alterations or improvement to real property)
- equipment that basically serves comfort or convenience functions (e.g., physical fitness equipment or trays, back packs)
- equipment that is primarily for the convenience a person caring for the member (e.g., cushion lift chairs)
- self help devices (e.g., safety grab bars)
- Power wheelchairs and scooters are not a covered benefit of CMS.

Rental versus Purchase

- The decision to rent or purchase DME will be made by CMS. All DME must be obtained from a vendor that accepts CMS pricing. If it can be determined that the equipment can be rented for a cost less than purchase payment then the rental will be made. Purchase may be covered only after it is proven through documentation that either:
 - the equipment is unobtainable on a rental basis, or
 - the patient will use the equipment for a long enough period of time to make its purchase more economical than continuing rental fees.
- Generally pieces of equipment such as wheelchairs, hospital beds, and oxygen have been rented, and walkers, bedside commodes, and bath benches have been purchased.

Repair or Replacement

 If a DME item requires repair or replacement, refer to the DME vendor contract or service agreement.

DME (DURABLE MEDICAL EQUIPMENT) (Continued)

 Repair of DME purchased by the patient may be covered if this DME item is a covered benefit of CMS. Repair of rental DME is not covered by CMS but may be covered as part of the contract or service agreement with the DME vendor.

Replacement of DME is covered by CMS only if the item is:

- medically necessary
- o ordered or prescribed by a CMS provider
- o provided by a vendor approved by CMS
- Replacement is not required because of misuse or loss by member.
- The following list includes some covered DME items:
 - o Blood glucose monitors
 - Canes (except white canes for the blind)
 - Commode chairs
 - o Crutches
 - "Grabbers" or other tools needed for a patient to be independent with ADLs
 - Home oxygen equipment and supplies
 - Infusion pumps (for IV antibiotics or Enteral Feedings)
 - Nebulizers
 - Walkers
- Prosthetic devices:
 - o Arm, leg, back, and neck braces
 - o Artificial limbs and eyes
 - Breast prostheses (including a surgical brassiere) after a mastectomy
 - o Ostomy supplies
- The following list includes DME items that should be delegated to a RN (not an AC) to evaluate:
 - C-Pap, Bi-Pap machines and supplies (purchase)

ME (DURABLE MEDICAL EQUIPMENT) (Continued)

- CPT machines (Post TKR) are rarely approved if a vendor accepts the reimbursement rate
- Dyna-Splints (patient should have completed a full course of PT –usually 8 sessions or 1 month- AND should be completing the recommended home exercises.)
- o Hospital beds
- Lymphedema pumps/pneumatic compression devices are rarely approved.
 Lymphedema clinic is the preferred choice.
- o Replacement of DME items.
- o TENS Unit and supplies.
- Wheelchairs Manual (RN discretion to rent or purchase, based on the diagnosis and estimated length of need)
- Wound Vacs (care taken to evaluate Diagnostic Codes and Dates of Service)

DUPUYTREN'S CONTRACTURE

Criteria for Authorization

Referral for this condition is only approved in cases in which the disease interferes with work or activities of daily living.

Patient History (both must be present)

- involvement of the palmar and digital fascia
- flexion deformity of the fingers

AND

Physical Exam (both must be present)

- characteristic nodule or cord in the palmar fascia
- metacarpophalangeal joint contracture
 >30 degrees

AND

Work History

ESOPHOGASTRODUODENOSCOPY (EGD)

Criteria for Authorization

EGD is also known as Upper GI Endoscopy and is performed when direct visualization of the upper GI tract is necessary. Biopsy of the esophagus, stomach and duodenum can also be done by EGD.

Patient History

- persistent symptoms of heartburn or GERD despite 2 months of PPI therapy.
- age of onset of GERD age 50 or later.
- extra esophageal symptoms, e.g., hoarseness, chest pain, wheezing
- complicated GERD, e.g. dysphagia or iron deficiency anemia
- symptoms of five years duration in patient > 50 years of age
- failure of lifestyle modifications such as no smoking, caffeine, aspirin, alcohol and spices

Repeat EGD (may be done in 8–12 weeks)

erosive or transitional cells present on initial biopsy

ENDOMETRIAL ABLATION

Criteria for Authorization

Endometrial ablation is an alternative to hysterectomy for women with persistent excessive vaginal bleeding. It is also used for women with hypertrophy or polyps of the endometrial tissue. It has the advantage over hysterectomy in that it does not require major surgery and preserves the uterus.

Patient History (all should be present)

- excessive vaginal bleeding in a woman who has completed childbearing
 - profuse bleeding or repetitive periods
 - anemia due to acute or chronic blood loss
- no uterine or cervical pathology that would require hysterectomy
- no finding of remedial cause by hysteroscopy
- failure of hormone treatment

EPIDURAL STEROID INJECTION

Criteria for Authorization

Epidural steroid injection is indicated for chronic neck or back pain with radiculopathy. It is an alternative to surgery, and may reduce the need for pain medications. Approval is given for only one injection at a time.

Patient History

- chronic neck or back pain with radicular symptoms present for at least 3 months.
- conservative pain management has been used for at least 6 weeks without benefit

Physical Exam

evidence of neurologic signs (numbness, weakness or reflex changes)

Addendum: Injections limited to three in a given year, with at least 3 months separation between injections.

EPILEPSY (SEIZURE DISORDER) VAGUS NERVE STIMULATION (VNS)

Criteria for Authorization

All patients with a current active seizure disorder should be seen by a neurologist at least once a year, even if seizure free. The number and types of seizure medications is changing frequently. Only a neurologist should remove seizure medication therapy. In those patients where medications are not effective for control, VNS or epilepsy surgery may be indicated.

Patient History (for VNS)

- failure of at least three drugs as therapy to control seizures
 - exclude discontinuance due to side effects and non-compliance.
- patient has the ability to manage a VNS magnet

ERYTHROPOIETIN (RECOMBINANT GROWTH FACTOR), EPOGEN, PROCRIT

Criteria for Authorization

The use of these medications has recently come under criticism for their failure to improve patient outcomes and their great expense. Medicare and other health insurance plans are increasing their restrictions on these medications. Erythropoietin is covered only for patients with severe anemia due to chronic kidney disease or cancer therapy to avoid blood transfusions.

Patient History (anemia criteria must be met for either condition)

- recurring chemotherapy expected to cause bone marrow suppression
- chronic kidney disease
- anemia with hemoglobin (Hgb) less than 10 gm/dl or hematocrit (Hct) less than 30%

Addendum:

- Dosages are approved for one month at a time, and are continued only if Anemia criteria are still present.
- Continuation of therapy for more than 6-8 weeks is not beneficial in the absence of response (e.g.<1 to 2 gm/dl rise in Hgb). Longer term use of the medications will not be approved in the absence of a response.

GANGLION CYST

Criteria for Authorization

Most ganglion cysts are painless and do not interfere with work or living activities. Approval for referral for surgery is limited to those patients who have a critical medical indication for surgery.

Patient History

- pain which causes interference with work or essential activities
- weakness or altered range of motion

Physical Exam

- cyst or mass of dorsal or volar wrist
- cyst or mass in other location causing a limitation of function

Addendum:

- 50% of ganglion cysts disappear without therapy
- regardless of therapy, reoccurrence is common

GENETIC COUNSELING

Genetic testing, treatment, or counseling begins with a consultation for Genetic Counseling. CMS covers Genetic Counseling only under the condition that Genetic testing of the member may change or inform the member's treatment plan.

Criteria for Authorization

Patient History (one of the three)

- Cancer that may be linked to a genetic predilection to other types of cancer, such as Hereditary Breast and Ovarian Cancer.
- A disease in which the diagnosis, severity or manifestation of the disease may be influenced by the genetic typing. For example, Multiple Endocrine Neoplasia type I or II.
- A disease in which the recommendations for treatment may depend on genetic typing. For example, Familial Adenomatous Polyposis.

CMS does not cover Genetic Testing, Treatment or Counseling for the following:

- Non-medical reasons (e.g., court-ordered tests, work-related tests, paternity tests).
- Non-medically necessary screening to determine carrier status for inheritable disorders when there would not be an immediate medical benefit or when results would not be used to initiate medical interventions/treatment.
- Testing of persons who have no clinical evidence or family history of a genetic abnormality.

GYNECOMASTIA

Abnormal enlargement of the breast in a male is usually due to hormonal imbalance or medications, especially anabolic steroids. The management of Gynecomastia is usually medical and a referral to endocrinology is indicated if the cause is unclear in the primary care setting. The basic work-up should be done in primary care, such as a careful medication history and hormone levels. CMS does not cover surgery for Gynecomastia unless there is a malignancy.

Patient History (must be documented)

- duration of gynecomastia
- medication history
- use of alcohol
- mammogram report (if done)
- significant weight gain
- history of liver disease
- onset of puberty

AND

Physical Exam

- unilateral or bilateral
- increased adipose vs. breast tissue

HEADACHE - MIGRAINES

Criteria for Authorization

The PCP is able to evaluate and manage most patients with migraines. Referral to a neurologist is only approved for a failure to respond to treatment or positive neurologic findings.

Patient History

- dissimilar headache (One item)
 - decreased alertness
 - first headache after age 50
 - "worst headache ever"
 - headache with exertion
- failure of adequate trial of management (Two of five)
 - nonsteroidal anti-inflammatories
 - isometheptine
 - eraotamine
 - butalbital
 - triptans (see below)

Physical Exam

- neurologic exam performed, any focal abnormality
- nuchal rigidity
- abnormal vital signs

Criteria for the approval of a triptin medication (e.g. Imitrex)

- requested by a Neurologist
- requested by PCP:
 - failed a non-steroidal medication
 - failed other generic anti-migraine medications

HEADACHE

Most patients with headache, including migraine, are managed by primary care. The following are criteria which may lead to a referral to a neurologist or other headache specialist. Advanced imaging (CT or MRI) is only indicated for headache when there are neurologic signs or an abnormal neurologic exam. CT is generally done in the emergency setting to rule out hemorrhage or mass lesion. MRI is preferred to evaluate for brain tumors and other intracranial lesions.

Note: A separate medical policy follows for Migraine.

Patient History

- sudden onset of new severe headache
- progressively worsening headaches
- onset with exertion, coughing, straining, and/or sexual activity
- associated symptoms such as:
 - drowsiness, confusion, memory loss
 - chronic malaise, myalgia, arthralgia
 - fever
 - progressive visual disturbances
 - weakness, clumsiness, loss of balance
 - onset of first headache after the age of 50 years

Physical Exam

- abnormal vital signs, especially fever or high blood pressure
- altered consciousness or cognition
- meningeal irritation ('stiff neck')
- papilloedema or fundal hemorrhage
- pupils unequal and/or poorly reactive
- weakness or sensory loss in face or limbs
- reflex asymmetry or abnormal plantar response
- clumsiness or loss of balance
- tender temporal arteries with diminished pulse

HEARING LOSS

The CMS program covers referral, testing and treatment for hearing loss which impairs a person's ability to work and handle activities of daily living. A work history is usually required for any treatment.

<u>Criteria for Authorization</u>

Patient History (At least one of these is required)

- infection or trauma suggesting a Conductive hearing loss. With infection (otitis media), the hearing loss must be present for at least 2 months
- otosclerosis
- sensorineural hearing loss with:
 - lesion of cochlea
 - tinnitus
 - gait imbalance
 - unilateral hearing loss
- a family history of a genetic hearing loss
- sudden onset of a major hearing loss
- recurrent dizziness with hearing loss
- poor speech discrimination

Physical Exam (At least one of these is present)

- otoscope Exam
 - no presence of blood, pus, cerumen plug, or foreign objects (all of which are treated in primary care)
 - abnormal findings of the tympanic membrane or middle ear which suggest a permanent or chronic problem

Tests

- audiogram shows evidence of more than a 30 decibel deficit
- replacement or repair-1x per 12 month period
- bi-aural hearing aids require visual acuity justification

HEMORRHOIDECTOMY

Criteria for Authorization

Most thrombosed external hemorrhoids resolve with warm baths, topical creams and fiber in the diet. CMS does not cover referral for treatment of external hemorrhoids unless the following criteria are met. Internal hemorrhoids which demonstrate recurrent bleeding and/or prolapse may warrant a procedure if they interfere with work or daily activities.

Patient History (any one of these present)

- repeated or persistent prolapse or thrombosis with severe pain (internal hemorrhoids)
- recurrent bleeding unresponsive to conservative treatment (either external or internal hemorrhoids)
- thrombosis with severe pain not responsive to warm baths or medications over 3 days

Physical Exam

acute irreducible prolapse of internal hemorrhoids

HEPATITIS B & C

Criteria for Authorization

A large number of CMS patients are Hepatitis B or C antibody positive. Of these patients, 55-85% will be chronically infected. 80-95% of patients with a positive Hepatitis C antibody will not develop serious liver disease such as cirrhosis or liver failure. The CMS program will only approve for referral and treatment those patients who are candidates for treatment.

In order to use limited CMS funds wisely for this population, only patients with some evidence of active liver disease from Hepatitis B or C will be referred for treatment. Referral requests should include: Hepatitis B or C antibody test, liver function tests (especially ALT) and information about other possible causes of liver disease such as active alcoholism, hyperlipidemia, weight and whether there is ongoing IV drug use.

For those patients who are Hepatitis B or C antibody positive, who have elevated liver function tests, and do not have another apparent cause for liver disease, HBV or HCV RNA testing should be done to evaluate for chronic infection. If the HCV RNA testing is positive, and the ALT is at least 50% above normal due to no other cause, a referral to GI for consideration for treatment will be approved. Also, if the HCV RNA testing is positive, HCV Genotype testing may be done by your clinic (if required by the GI referral specialist) or by the specialty clinic.

Patients who are Hepatitis B or C antibody positive and have normal liver function tests should be followed clinically at the primary care level. Also, patients with ongoing alcoholism, IV drug use, fatty liver or other causes of liver disease should have these conditions managed before consideration for hepatitis C referral for treatment.

This link to the CDC website office recommended Clinical Guidelines for the Diagnosis and Treatment of Hepatitis:

http://www.cdc.gov

https://www.aasld.org

HERNIAS - SURGICAL CORRECTION

<u>Criteria for Authorization</u>

The CMS program does not cover elective surgery for hernia repair. Current clinical guidelines support "watchful waiting" for hernias in patients who are not at high risk for incarceration. The near-elderly and those persons doing heavy manual labor are at risk. The CMS program will only cover for referral and treatment of hernias that are symptomatic and interfere with work or activities of daily living.

Inguinal, Incisional, Ventral or Peri-umbilical Hernias:

Patient History (one of two)

- pain of significant duration
- affects employability

AND

Physical Exam (one of three)

- difficulty reducing hernia
- incarcerated hernia
- tenderness upon examination

AND

Work History

HIRSUTISM

Criteria for Authorization

Hirsutism is the excessive growth of hair in women. A family and cultural history are important to be sure that the hair growth is not normal. People from Mediterranean countries often have excessive hair growth including some masculinization of women. Hirsutism is triggered by androgen production, and the work-up focuses on hormonal causes. The work-up can usually be done in primary care, but based on the criteria below, a referral to an endocrinologist, or an gynecologist may be appropriate.

Patient History (all are present)

- symptoms suggestive of polycystic ovarian disease or adrenal hyperplasia, Hyperandrogenic, insulinresistant, acanthosis nigricans syndrome and androgen secreting tumors
- absent familial predilection for hirsutism
- evaluation of patient's medications

AND

Physical Exam (both are done)

- confirmation of hirsutism
- pelvic ultrasound

HORMONE THERAPY

Criteria for Authorization

Oral Contraceptives Pills (OCPs): The CMS program does not cover contraceptive care. OCPs are only approved for specific medical indications:

Patient History

- dysfunctional bleeding
- polycystic ovary syndrome
- hypothalamic amenorrhea
- endometriosis
- recurrent functional ovarian cysts

Progesterone Challenge (usually 10-14 days)

As an alternative to surgery or prevention of surgery

- dysfunctional bleeding
- secondary amenorrhea
- Hemoglobin < 10 mg/dl

Lupron

Prescribed only as recommended by a gynecologist

- fibroids
- endometriosis
- limited to 3 mg (1x per month)

Hormone Replacement Therapy (HRT): CMS does not cover HRT for normal menopause

May be prescribed for:

- surgical menopause < 40 years of age (include date of surgery), and approve only to age 50
- osteoporosis, documented by Dexa Scan
- progesterone used with estrogen if uterus present
- other congenital estrogen deficiency

HYDROCELE

<u>Criteria for Authorization</u>

Hydrocele is rarely a significant health problem and CMS does not cover routine referral or correction.

Patient History

- painless mass
- vague, gradual symptoms may occur with enlargement

Physical Exam

- mass or focal swelling
- cystic
- freely movable
- non-tender
- transluminates with light
- there may be testicular atrophy
- usually left-sided

Diagnosis

 based on translumination with a lack of any mass or solid tissue. Ultrasound may be performed if exam is uncertain

Management

- observe for spontaneous resolution or if any change
- no treatment necessary if stable
- aspiration may be performed if enlarging, causing discomfort or interfering with work
- criteria for surgical removal
 - persistent pain
 - interferes with work

HYPERPARATHYROIDISM (CAUSING HYPOCALCAEMIA)

Criteria for Authorization

Patient History and Laboratory Findings:

- occurrence of renal stones
- progressive bone loss (by Dexa Scan)
- serum CA > 11.5 mg/d or simultaneous elevation of serum PTH and Calcium indicating hyperparathyroidism

HYPOGONADISM AND TESTOSTERONE REPLACEMENT

CMS only approves testosterone replacement for therapeutic treatment of a medical condition. Hypogonadism is defined as "inadequate gonadal function, as manifested by deficiencies in gametogeneisis and/or the secretion of gonadal hormones". Hypogonadism may manifest with testosterone deficiency, infertility, or both conditions. Symptoms of hypogonadism depend primarily on the age of the male member at the time of the development of the condition.

CMS does not provide coverage for treatment for members with a diagnosis of infertility, male erectile dysfunction and/or loss of or decrease of libido under any circumstances.

CMS may cover treatment for members with conditions of primary hypogonadism, such as Klinefelter syndrome, Mumps Orchitis or Hemochromatosis, or with conditions of secondary hypogonadism, such as Kallman syndrome, Inflammatory disease, or Pituitary disorders.

Criteria for Authorization

Hormone replacement with testosterone preparations for hypogonadism (excluding injectables) may be considered under the following circumstances:

- Treatment of AIDS Wasting Syndrome
- Osteopenia or osteoporosis as confirmed by Bone Density Study associated with hypogonadism that develops postpubertal as a result of one of the following:
 - o genetic syndromes
 - o congenital disorders
 - o toxic exposures
 - radiation treatment
 - o chemotherapy
 - o cryptochidism
 - o mumps orchitis
 - o hemochromatosis
 - o autoimmune syndromes
 - testicular trauma
 - o secondary to hypothalamic-pituitary tumors

HYPOGONADISM AND TESTOSTERONE REPLACEMENT (Continued)

- For post-pubertal hypogonadism testosterone replacement may be considered when the member has:
 - two (2) morning laboratory tests confirming low total testosterone level of less than 200ng/dL on each test.
 - o AND Hypogonadal symptoms
- Testosterone replacement therapy is contraindicated in men with prostate cancer, male breast cancer, or untreated prolactinoma.
 Sleep apnea and polycythemia, which may cause hyperviscosity, are relative contraindications to the use of testosterone therapy.

HYSTERECTOMY

Criteria for Authorization

The CMS program does not cover elective hysterectomy. The procedure is only approved when critically necessary for the woman's health. Abdominal, vaginal or laparoscopic hysterectomy is only approved when clinically necessary. Less invasive procedures, such as uterine ablation therapy should always be considered.

Patient History

- cancer or pre-cancer of the uterus, cervix or ovary
- recurrent endometrial hyperplasia after adequate treatment with curettage and progestin therapy
- rapid growth of fibroids which are causing health problems such as persistent heavy vaginal bleeding
- progressive dysmenorrhea or menorrhagia unresponsive to D&C, hysteroscopy and hormone therapy
- refractory menorrhagia for 3-6 months despite adequate hormone therapy. There must be a clinically significant drop in Hb or Hct.

OR

Physical Exam

 the presence of uterine fibroids > 16 wks gestational size

AND

Hysterectomy worksheet required

REQUEST FOR AUTHORIZATION HYSTERECTOMY WORKSHEET

<u>History</u>			
Age	Gravida	Parity	
Symptoms: •	Recurrent Bleeding Duration Lowest Hb/Hct (with documentation) Current Hb/Hct*		
<u>Treatment</u> • • • •	Oral Contraceptives (3 of Depo-Provera (3 cycles) D&C or Hysteroscopy* Intractable Pelvic Pain	•	
Duration	Cyclic Constant Treatment Medications Duration Presence of Fibroids Size >16 weeks*		
•	Pelvic Relaxation - Severi - Uterine Prolapse - Cystocele - Rectocele Pelvic Tenderness Abnormal Findings	ty	
*provide do	ocumentation		
Patient Name:		DOB:	
Date completed		Complet	ed By:

HYSTEROSCOPY

Criteria for Authorization

Hysteroscopy is an outpatient procedure allowing the physician to visualize the inside of the uterus. Hysteroscopy is superior to D&C for diagnosis of intrauterine pathology. Endometrial biopsies and endometrial ablation may be done as part of Hysteroscopy.

Patient History

- postmenopausal bleeding
- failure to find cervical or uterine pathology that would cause abnormal bleeding
- history of excessive uterine bleeding evidenced by profuse bleeding, repetitive periods lasting more than 8 days or frequent periods at less than 21-day intervals
- failure of appropriate medical therapy

AND

Physical exam and Diagnostics

- pelvic exam
- obtain cervical cytology
- obtain endometrial sampling

INCONTINENCE

Criteria for Authorization

Most urinary incontinence in women is managed by primary care with pelvic exercises and medication. Referral for surgical correction is only approved if critically necessary for employment or daily activities.

Patient History

- duration of symptoms
- thoroughly evaluated and treated with behavioral techniques and medication
- patient has previously followed a mandatory voiding schedule with specific fluid intake, i.e. no caffeinated beverages
- compliance with a pelvic exercise program (Kegels)
- trial of medications for incontinence
- alternative use of pessary offered to patient

AND

Physical Exam

- assessment of estrogen status (evidence of atrophy)
- adequate pelvic exam, R/O diverticula and fistulas, description of prolapse
- urine culture

INFLUENZA (INCLUDING SWINE FLU)

The CMS formulary does not cover anti-viral agents for influenza (the only anti-viral agent covered is acyclovir for herpes).

The generic anti-viral agent (amantadine) and Flumadine are not effective against the swine flu.

Tamiflu and Relenza are expensive and only recommended in highly suspect patients that are either seriously ill or if being treated in the first 48 hours.

Use of the anti-viral drugs should be a code 1 authorization request.

The most current recommendation of the CDC and the San Diego County Public Health Department will be followed.

INSULIN PUMP

Criteria for Authorization

- request from Endocrinologist
- patient must have frequent and severe glycemic events requiring visits to Physician, ER or Hospital.
- significant ketosis
- insulin reactions and/or ketoacidosis
- blood glucose levels greater than 140 mg/dL preprandially and/or greater than 200 mg/dL fasting ("Dawn phenomenon")
- glycosylated hemoglobin (HbA1c) greater than 8 percent
- chronic renal failure or ongoing dialysis
- intermittent insulin injection not a practical option for the patient

Note: If insulin pump in place and patient is doing well, CMS will provide supplies to maintain it.

LAMINECTOMY (SPINE SURGERY)

Criteria for Authorization

Surgery to the spine, cervical, thoracic or lumbar, is done for nerve impingement not responsive to conservative measures. Physical therapy and epidural steroid injections should be tried first in most cases.

Patient History

- radiating pain from lumbar spine down leg
- numbness of leg or foot
- low back pain
- bowel or bladder dysfunction

AND

Physical Exam

- ↓ sensation
- + contra lateral straight leg raising
- ↓ DTR (deep tendon reflexes)
- ↓ muscle strength
- change in gait
- unequal deep tendon reflexes

AND

Diagnostics

MRI is the imaging of choice. A CT scan may be adequate if already done

- demonstrates positive disc protrusion
- spinal stenosis
- cord compression with neurological sign

LAPAROSCOPY (GYNECOLOGY)

<u>Criteria for Authorization</u>

- chronic pelvic pain with no cause identified
- abnormal ovarian findings
- failure of conservative management (OCs, progesterone)
- failure of GnRH Agonist (endometriosis)

MAMMOGRAPHY

Criteria for Authorization

The CMS program does not cover routine screening mammography. A diagnostic mammogram is only approved when critically necessary to evaluate an abnormality suggesting possible breast cancer.

• If the woman is eligible, refer to the Breast and Cervical Cancer Early Detection Program (BCCEDP).

MEDICAL TRANSPORTATION

Criteria for Authorization

Medical transportation for certified CMS patients is coordinated in the Medical Management Services Department. There are three categories of medical transportation: Emergent (by ambulance with an ACLS certified team of EMTs), Urgent (transport with a BLS trained team), and Non-Urgent medical transportation (wheelchair van, transport on a stretcher/gurney).

Emergent transportation for medical care (must be documented)

- Ambulance transport to the emergency department when medically necessary (Patient requires the care of a certified EMT during transportation, and/or requires immediate medical attention that can not be obtained by other means.)
- Ambulance transport from one hospital to another to facilitate the prompt receipt of appropriate services, as medically necessary.

Urgent transportation for medical care

- Patient does not require the supervision and services of an EMT or ACLS certified individual during transport.
- Non-emergent medical transport to the Emergency Department for a patient who has been stabilized by the Primary Physician.

Non-Urgent transportation for medical care

- Patient does not require medical supervision during transportation.
- Patient's medical condition makes it unsafe or impossible to be transported in other type of vehicle. (For example, patients who have been casted or fixated in a position that requires a stretcher, patients

MEDICAL TRANSPORATION (Continued)

requiring wheel chair transportation if they are unable to transfer independently from their wheelchair.)

Transportation upon hospital discharge

- The acute care facility is responsible for transporting the patient upon discharge, as needed.
- CMS does not cover transportation to facilities that are not covered under CMS scope of service (mental health institutions, drug or alcohol rehabilitation services).

MRI

Criteria for Authorization

An MRI should only be performed if the diagnosis or extent of the disease is unknown. Written progress notes and an order must be submitted in order to document the need for the imaging. The MRI is used most commonly to image soft tissue to look for disease not visible on regular x-rays. In most cases, plain x-rays are done first and if common osteoarthritis is found an MRI is not necessary. An MRI of the spine requires a history of persistent radicular symptoms and findings on a neurologic exam that confirm nerve entrapment. An MRI of the brain requires positive neurologic findings on physical exam. For other areas of the body, suspected malignancy, suspect osteomyelitis and significant interference with work or activities of daily living are important criteria for approval. An MRI should only be done when its findings will have a direct result on the diagnosis and treatment of the patient.

Milliman Ambulatory Care Guidelines should be consulted if there is any question about the use of MRI to various areas of the body.

MRI OF KNEE

Criteria for Authorization

A MRI of the knee should only be performed if the diagnosis or extent of the disease is unknown. Plain x-rays of the knees are done first and if common osteoarthritis is found, a MRI is not necessary. A careful knee exam should also be performed before consideration is given for an MRI. A MRI of the knee should be performed before an Orthopedic consultation for most knee problems.

Patient History

- aid in the diagnosis of meniscal tear
- aid in the diagnosis of an internal ligament tear
- detection, staging, post-treatment evaluation of tumor of the knee
- suspected osteochondritis dessicans if the clinical picture and plain x-rays are not confirmatory
- suspected osteonecrosis if the clinical picture and plain x-rays are not confirmatory
- persistent knee pain/swelling and/or instability
 after an injury which has not responded to
 conservative management (ice, rest, elevation,
 medication, non-weight bearing, physical
 therapy), if plain x-rays have failed to
 demonstrate a fracture or loose body, and if the
 clinical picture is unclear
- persistent knee/pain swelling and/or instability (gives way) not associated with an injury after a 3-6 week trial of conservative treatment
- if specifically requested by a consulting physician (orthopedist or rheumatologist)

Addendum: An MRI is not indicated for:

- Diagnosis of osteoarthritis or rheumatoid arthritis.
- Diagnosis of torn meniscus, loose body, or osteochondritis dessicans when the clinical examination and x-rays are diagnostic. If there is a true "locking" of the knee in flexion rather than

MRI OF KNEE (Continued)

- "catching" in extension, this is indicative of loose body or torn meniscus.
- When the MRI results will not alter the treatment plan of an anticipated surgical procedure.

NEPHROLOGY

Criteria for Authorization

Referral to Nephrology is most often considered for patients with renal failure. Other indications include chronic renal stones and other renal disease. In patients with hypertension and diabetes, monitoring renal function is vitally important, since these diseases are the most common causes of renal failure.

Mild renal insufficiency is managed at the primary care level. Counseling and appropriate adjustment of medications with declining renal function are important management consideration. Referral to Nephrology and follow-up is approved for patients with a serum creatinine of 2.0 or higher or if the estimated GFR is 30 or lower.

OCULAR DISEASE - REFERRAL FOR VISION LOSS OR EYE PAIN

Criteria for Authorization

The CMS program does not cover routine eye care, including refractions. The program will cover critical eye services necessary to allow a patient to work and to relieve pain. Treatment Authorizations Requests for refraction are covered only for patients experiencing vision loss to the extent that it interferes with work and basic life functions. All conjunctivitis is treated by primary care.

Patient History (one of three)

- decreased visual acuity (provide visual acuity)
- ocular pain
- photophobia

Physical Exam (one of three must be present unless vision loss is documented)

- injection of vessels around the cornea
- corneal opacification
- pupil abnormalities
- suspicious for corneal ulceration

OPHTHALMOLOGY

Chalazion (is a cyst in the eyelid that is caused by inflammation of the meibomian gland).

The primary treatment is application of warm compresses for 10 - 20 minutes at least 4 times a day. This may soften the hardened oils blocking the duct and promote drainage and healing.

Topical antibiotic drops or ointment are sometimes used for the initial acute infection, but are otherwise of little value in treating a chalazion. Chalazia will often disappear without further treatment within a few months and virtually all will reabsorb within two years.

If they continue to enlarge or fail to settle within a few months, then a referral to an ophthalmologist is appropriate. Smaller lesions may be injected with a corticosteroid or larger ones may be surgically removed using local anesthesia.

Criteria for Authorization

Patient History

persistent lesion (3 months or longer)

Failure of Treatment

 conservative therapy with antibiotic and warm compresses for two months and/or local injection of a corticosteroid

Blepharitis (inflammation of the eyelids)

Many forms of treatment will improve blepharitis, including both antibiotic or steroid eye drops, and certain oral antibiotics. Unfortunately it may recur when any treatment is ceased. Recommend a regime of daily eyelid cleaning which is both effective and can be continued safely long-term. Simply cleaning the eyelids with a face cloth during every bath or shower may be a good system for a patient.

OPHTHALMOLOGY (Continued)

Patient History

- failure of improvement despite treatment Physical Exam
 - Persistent Inflammation of the lid margins

<u>Iritis (or Uveitis)</u> Iritis is inflammation predominantly located in the iris of the eye. Inflammation in the iris is more correctly classified as anterior uveitis. The ciliary body can also be inflamed and this would then be called iridocyclitis.

Criteria for Authorization

Patient History (all three required if no physical findings present)

- photophobia
- moderate pain
- vision is blurred

Physical Exam (any one item)

- redness of the sclera
- red halo around the cornea
- a clear discharge may be present

OPHTHALMOLOGY - CATARACTS

Cataract: An opacity that develops in the crystalline lens of the eye or in its envelope.

<u>Criteria for Authorization</u>

Patient History

blurred vision (provide visual acuity)

Physical Exam

 cornea is clouded and cataract is seen on fundus exam

Criteria for Surgical Removal

History (both required)

- functional impairment employment and/or ADL's affected
- failure of vision to improve with prescription changes and/or other corrective measures

AND

Physical Exam

Visual acuity in best eye must be worse than 20/50 with corrective lenses. (Surgery is covered for both eyes.)

Addendum: A cataract may be removed at any level of acuity if it precludes diagnosis or treatment of another ocular disease, such as diabetes or natural disease.

OPHTHALMOLOGY - GLAUCOMA

Criteria for Authorization

Patient History

- loss of the mid-peripheral visual field
- elevated intraocular pressure
- advanced age
- African-American ethnicity
- family history of glaucoma
- other risk factors
 - myopia
 - diabetes mellitus
 - migraine
 - hypertension
 - long-term corticosteroid use
 - previous eye injury

Physical Exam

suspicious looking optic nerve head

Addendum: According to A.A.O. (American Academy of Ophthalmology).

- > 60 years exam every 2 years
- > 40 years in Blacks
- 20-39 years in Blacks exam every 3-5 years

OPHTHALMOLOGY - PTERYGIUM

Criteria for Authorization

Pterygium is fibrous material that forms in the eye and covers part of the cornea. This benign condition often occurs in persons chronically exposed to dust and outdoor conditions. Referral for surgery is only necessary when vision is impaired.

Patient History

• visual interference (provide documentation)

AND

Physical Exam

 extension onto or over cornea to the extent that vision is impaired

OPHTHALMOLOGY - FOLLOW-UP EXAMINATION GUIDELINES

<u>DIAGNOSIS</u>: **ANTERIOR ISCHEMIC OPTIC NEUROPATHY** – every 2 weeks

for 2 visits, then every 3-6 months

DIAGNOSIS: **BLEPHARITIS** – up to 2 visits annually

DIAGNOSIS: CATARACT - If immature cataract, every 12 months.

If post-operative – covered in global, every 3 months. If post capsule thickening, every 6 months. If best

corrected visual acuity is 20/40 or worse in the best eye, every 6 months. If following surgery in eye #1, when best corrected visual acuity is 20/40 in the remaining eye,

every 6 months.

<u>DIAGNOSIS</u>: **CHLOROQUINE RETINOPATHY** – every 6-12 months

<u>DIAGNOSIS</u>: **CORNEAL ABRASION** – every 1-2 days until healed

<u>DIAGNOSIS</u>: **CORNEAL ULCER** – every 24 hours until healed

<u>DIAGNOSIS</u>: **CYSTOID MACULAR EDEMA (CME)** – every 6 weeks to 3

months, depending on medication used.

DIAGNOSIS: **DIABETES MELLITUS** (DM), - annually for retina exam

With retinopathy – every 3-6 months

<u>DIAGNOSIS</u>: **DIABETIC MACULAR EDEMA** – every 3 months

DIAGNOSIS: **EPIRETINAL MEMBRANE (ERM)** – every 3 months

DIAGNOSIS: GIANT CELL ARTERITIS (VASCULITIS) – as often as needed

based on the stability of the patient and nature of steroid

therapy

<u>DIAGNOSIS</u>: **GLAUCOMA (chronic)** – Every 3 months

DIAGNOSIS: GLAUCOMA SUSPECT – every 3-6 months depending on

the pressure

DIAGNOSIS: **HYPERTENSIVE RETINOPATHY** – every 6-12 months

OPHTHALMOLOGY - FOLLOW-UP EXAMINATION GUIDELINES (Continued)

<u>DIAGNOSIS</u>: **HYPERTENSIVE RETINOPATHY** – every 6-12 months

<u>DIAGNOSIS</u>: **IRITIS, UVEITIS, IRIDOCYCLITIS** – every 1-2 weeks if acute,

every 1-3 months if chronic depending on medication

used and severity of the inflammation

<u>DIAGNOSIS</u>: **MACULAR DEGENERATION** – every 1-12 months

depending on severity and progression of the disease

DIAGNOSIS: MACULAR HOLE – every 1-3 months

DIAGNOSIS: **OPACIFICATION OF POSTERIOR CAPSULE** – once a year

DIAGNOSIS: OPTIC NEURITIS – as often as needed

DIAGNOSIS: **PSEUDOPHAKIA** – once a year

DIAGNOSIS: **RETINAL DETACHMENT** – as often as needed before or

after surgery

<u>DIAGNOSIS</u>: **RETINAL VEIN OCCLUSION (Central or Branch) -** Every

month for 3 months, then every 3 months until stable,

then every 6 months as needed

DIAGNOSIS: **RETINITIS PIGMENTOSA** – once a year

DIAGNOSIS: STEROID EYE DROP USE – once a year

DIAGNOSIS: STEROID SYSTEMIC MEDICATION – once a year

<u>DIAGNOSIS</u>: TAMOXIFEN RETINOPATHY – every 12 months, including

Visual Field

<u>DIAGNOSIS</u>: **VISUAL FIELD DEFECT** – once a year

DIAGNOSIS: VITREOUS DETACHMENT OR FLOATER – with symptoms

such as flashes of light, every 3-6 months, otherwise once

a year

DIAGNOSIS: VITREOUS HEMORRAGE – every 1-3 weeks, ultrasound as

needed

OPTOMETRY- COVERAGE FOR GLASSES

Criteria for Authorization

- vision defect by Snellen testing of equal to or > 20/50 or change in any meridian by at least 1.0 Diopter from the previous prescription
- correction required for employment

Changes in Prescription:

- any meridian change by at least 1.0 diopter
- astigmatic correction of .5 diopters or more

Replacement:

• replacement of glasses - 1x in 12 months if broken, lost or stolen

Not a Benefit:

- reading glasses
- routine refractions without any noticeable change in vision

ORGAN TRANSPLANTATION

CMS never covers organ and bone transplant services or services related to getting an organ transplant. CMS may cover services related to avoiding rejection and/or medical complications that arise from an organ transplant received when a patient was not on the CMS program. Any such coverage would be based on a review by medical management.

OTITIS MEDIA

Criteria for Authorization

Acute otitis media resolves most of the time in 2-8 weeks. Treatment is performed in primary care. Chronic serous otitis media refers to a persistent collection of fluid in the middle ear. This usually resolves spontaneously or is accompanied by allergic rhinitis. The following criteria are used for referrals to ENT:

Patient History (one present)

- 3 or more episodes of acute otitis media in 6 months
- persistent pain and pressure in the middle ear longer than 3 months

Physical exam (one present)

- visualization of the tympanic membrane shows bulging, retraction or fluid layer
- inflammation of the tympanic membrane
- reduced mobility testing of the tympanic membrane by:
 - tympanometry
 - acoustic reflectometry

Decreased hearing by audiometry

PAIN MANAGEMENT

Criteria for Authorization

Most patients with chronic pain are managed by primary care. Referral to a Pain Specialist is appropriate if the diagnosis of chronic pain is uncertain despite a work-up by primary care, if a procedure such as an epidural injection is indicated, or if specialist help is needed with pain management.

Patient History (at least 7 must be present in the clinical record)

- failure of adequate medication treatment, NSAIDS, opiates
- failure of an individualized proactive pain control plan
- failure of relaxation exercises
- failure of a home exercise program
- patient has knowledge of and expectation of his or her pain management
- measurement of pain on a scale of 1-10
- failure of therapeutic modalities, heat, cold, physical therapy
- no evidence of drug addiction or drug-seeking behavior
- narcotic contract in place with clear boundaries and limitations
- documentation of previous surgeries

AND

Physical Exam (at least 1 present)

- restriction of movement
- sites of tenderness
- neurologic signs with neck or back pain, especially radiculopathy

AND

Tests done in primary care (Imaging and other studies) are required.

PEPTIC ULCER DISEASE

Criteria for Authorization

Peptic ulcer refers to gastric and duodenal ulcers. The most common causes are chronic h. pylori infection and the use of NSAIDs. Gastric ulcers carry an increased risk of cancer. Medical management by primary care is usually sufficient to treat peptic ulcers. Referral to GI is done for endoscopy (EGD). Because of the effectiveness of modern medical management, surgery for peptic ulcer disease is rarely necessary.

Patient History

- intractable and recurrent epigastric pain
- adequate trial of a PPI medication (at least 2 months therapy)
- treatment for h. pylori not successful

AND

Physical Exam

• epigastric tenderness is present.

AND

Labs

- h. pylori testing
- fecal occult blood

PET SCAN (POSITRON - EMISSION TOMOGRAPHY)

<u>Criteria for Authorization</u>

A PET scan differs from a CT or MRI by imaging cellular function of tissue. It is most useful in cancer diagnosis and follow-up. PET scanning is now being applied to other organ systems such as the heart. The CMS only covers PET scanning when it is critically necessary and no other modality will give the information.

- Diagnosis, staging and restaging or the following clinical conditions:
 - lung cancer (non-small-cell)
 - esophageal cancer
 - colorectal cancer
 - lymphoma
 - melanoma
 - head and neck malignancy
 - brain malignancy
 - thyroid cancer
 - breast cancer
 - cervical cancer
- To determine appropriate treatment, surgery v. chemotherapy.
- To determine if a tumor has been completely eradicated, post treatment.

PHYSICAL THERAPY, OCCUPATIONAL THERAPY, SPEECH THERAPY

Criteria for Authorization

PT, OT and ST are approved only for clinical conditions which require them for return to function. In general, an evaluation and 2 follow-up treatments are approved initially, unless the condition dictates longer therapy (such PT in a long bone fracture or OT/ST post stroke).

- ordered by the PCP when office education is not sufficient
- ordered by specialist
- post-op surgery
- must include aggressive patient education and home exercises
- greater than 12 patient visits require case management
- document action of progress and modalities used
- chronic low back pain unresponsive to medication management

Patient History must include a description of limitations Physical Exam must include deficits or impairment Patient Responsibility

- motivation to follow exercise program
- "No shows" and non-compliance will lose approval for further treatment

Addendum: Extension of therapy requests must be accompanied by original evaluation and comparative documentation to allow assessment of improvement in function.

PLANTAR FASCIITIS

Criteria for Authorization

Most patients with plantar fasciitis are managed by primary care. All patients should have their feet and shoes evaluated. Initial treatment includes heel lifts or shoe inserts. Decreased activity, stretching and weight loss are part of conservative treatment. Only persistent and severe cases are approved for referral.

Patient History

- Heel Pain:
 - increased upon awaking
 - located just anterior to the heel

AND

- Failure to respond to conservative management
 - modify activity
 - limit weight bearing (running)
 - oral anti-inflammatory medications
 - heel pads (OTC products)
 - hot soaks with no symptomatic relief
 - exercises to strengthen lower leg muscles and to increase flexibility of the achilles tendon and hamstrings

AND

Physical Exam

- pain to palpitation at plantar aspect of the calcaneous
- may have positive heel squeeze test
- often associated with pes cavus or pes planus

PODIATRY

Referral to Podiatry or an Orthopedist for foot problems is covered by CMS for conditions which are critically necessary for work and/or activities of daily living.

Most common foot problems such as plantar fasciitis and skin conditions are managed by primary care.

Proper shoes and the use of inserts treat most foot problems, along with weight loss and stretching.

Routine foot examinations, including the annual diabetic foot exam, are done by primary care.

CMS does not cover referral or treatment of toenail fungus infestation (tinea unguum).

<u>Criteria for Referral to a Podiatrist or an Orthopedist</u>

- major foot deformity, including a bunion which is causing pain and inability to work or perform ADLs
- failure of conservative care provided by PCP
- heel spurs failure to respond to conservative care and requiring an injection

PROSTATECTOMY

Criteria for Authorization

There are now multiple options for treating prostate disease other than surgery. Medications are very effective for benign prostatic hypertrophy (BPH). Surgery is only one of many options for treating prostate cancer. The decision to remove the prostate requires a specialist in Urology or Urologic Oncology.

Patient History (one of four)

- history of urinary retention despite treatment with medications
- presence of incontinence
- recurrent urinary infections
- nocturia more than twice/night despite treatment with medications

AND

Physical Exam (one of three)

- enlarged prostate or with malignant nodules
- > 20 c.c. post-voiding residual despite medications
- persistent hematuria

AND

Diagnostic (one of four)

- IVP obstructing prostate
- U/S hydronephrosis
- positive needle biopsy for cancer
- abnormal creatinine level, or decreased creatinine clearance

RHEUMATOLOGY - INCLUDING RHEUMATOID ARTHRITIS

Criteria for Authorization

Referral to Rheumatology must have evidence of active collagen vascular (autoimmune) disease. A thorough history, musculoskeletal exam and laboratory studies below must be done. The treatment of Rheumatoid Arthritis has changed to the early use of disease modifying medications to prevent further disability. Early referral to Rheumatology is now the standard of care. An accurate diagnosis of the condition should occur at the primary care level.

Patient History (three of six)

- chronic pain
- loss of joint function
- limitation of self-care (Noticeable joint inflammation, stiffness, deformity)
- morning stiffness, warmth, redness, swelling, and deformity
- any loss of finger function
- generalized illness

ACR – Diagnostic Criteria (three of seven)

- morning stiffness of at least 60 minutes duration
- arthritis of three or more joints
- arthritis of hand joints
- symmetric arthritis Present for six weeks
- rheumatoid nodules
- serum Rheumatoid Factor
- radiographic changes

Physical Exam

fusiform swelling of small joints, especially the hands

Laboratory (the first four must be provided)

- ANA (Antinuclear antibodies)
- Rheumatoid Factor
- ESR (Erythrocyte Sedimentation Rate)
- CBC, electrolytes, creatinine, liver function tests, UA
- synovial fluid analysis

SECOND OPINION

Criteria for Authorization

CMS may authorize a request for a second opinion from the patient or practitioner or CMS may suggest a second opinion when any one of the following circumstances are present:

- A more cost-effective treatment option is available.
- Conservative therapy has not been attempted or has not had sufficient time to show results.
- Practitioner or patient disagrees with the diagnosis and/or plan of treatment recommended by the specialist.
- Practitioner or patient is seeking an alternate treatment option that may improve the outcome.
- Patient/practitioner relationship is hindered.
- Geographic and/or other obstacles prohibit patient form accessing care.

SEPTOPLASTY

Criteria for Authorization

The CMS program does not cover surgery on the nose for any cosmetic purpose. Approval for referral and surgery would be approved only if the abnormality markedly interferes with work or daily function.

Patient History (one of two)

- documented nasal trauma of recent origin which causes significant nasal obstruction
- persistent serous otitis media secondary to nasal obstruction and Eustachian tube dysfunction

AND

Physical Exam

 dislocation of septal cartilage causing complete or near complete obstruction of the nasal cavity

SINUSITIS - ACUTE AND CHRONIC

Criteria for Authorization

Most acute and chronic sinusitis is managed by primary care. Underlying causes such as allergy and infection are determined and treated by primary care. Referral is only necessary for persistent disease that markedly interferes with work or daily function.

Patient History (one of two)

- persistent obstruction beyond two months which interferes with function
- failure to respond to 2-3 courses of antibiotic therapy

Physical Exam (one of two)

- nasal exam documenting purulent discharge
- palpable sinus tenderness

Treatment (three of three)

- first and second line antibiotics used for up to 6 weeks
- decongestant therapy for up to 6 weeks
- nasal corticosteroids used for up to 6 weeks without benefit

Diagnostic

• sinus x-rays or CT scan confirm obstruction

SLEEP APNEA

Criteria for Authorization

Sleep Apnea is common among obese people and people with upper airway obstruction. The sleep apnea form must be completed to help determine level of need. CPAP is the treatment of choice for documented clinically significant sleep apnea unless there is a removable obstructive lesion. Heat and humidification may be included with CPAP as standard therapy.

Patient History: (two of first four)

- chronic loud snoring
- gasping or choking episodes
- excessive daytime sleepiness
- cognitive difficulties
- stable home situation. (required)
- willingness to use C-PAP machine if recommended with electricity available in patient's bedroom (required)

Physical Exam:

- obesity, including nuchal obesity
- hypertension
- nasopharyngeal narrowing
- BMI > 35

Treatment:

- Oral Surgery only if there is an obstructive lesion that will relieve the problem.
- CPAP CPAP equipment is purchased from an approved DME provider. The usual life span for the equipment is 3-5 years. Replacement is on an as needed basis with a recommendation of the DME provider after an evaluation of the equipment.

COUNTY MEDICAL SERVICES STATEMENT OF MEDICAL NECESSITY REQUEST FOR SLEEP APNEA STUDY

Patient Name		_ SSN	Date	
CMS Eligible From:	To:	Clir	iic	
Requesting Practitioner		SSNDate Clinic Contact #		
Please complete the information Sleep Apnea Study.	n below to c	determine t	he medical necessity of a	
Patient History of Sleep Disturbat	<u>nce</u>			
Average number of hours of slee	ep each nig	ht	_	
Does patient nap during the da	ıy? □ Occa	sionally i	□ Daily	
Snoring: □Soft □ Loud				
□ Falls asleep while driving	□ Excessiv	e daytime	somnolence	
$\hfill \square$ Wakens with a sensation of ch	noking or ga	sping		
Medical Conditions				
□ Hypertension □Controlled □	Malignant	□ Allergic	Rhinitis	
□ Asthma □ Depression	□ Nocturio	a 🗆 Diabe	etes 🗆 Type I 🗆 Type II	
Controlled UYes UNo				
□ Obesity □ Heart Disease				
<u>Life Style Behaviors</u>				
Number of caffeinated beverag	ges per day			
Amount of alcohol consumed	□ Daily		ccasionally	
Smokes more than 1 pack of tol	bacco per c	lay 🗆 Yes 🗆	No	
Does the patient have a stable	home enviro	onment? 🗆	Yes □ No	
Medical Exam (all required)				
Height Weight B	lood Pressur	e N	eck circumference	
Adeno-tonsillar enlargement	□Y	es i	□No	
Maxillo-mandibular malformatic			□No	
Medications (list all)				
The practitioner has discussed th	ne treatmen	t options w	ith the patient. \square Yes \square No	
If a CPAP is indicated, the patie treatment (equipment, noise, dr	_		he inconvenience of the	
If a CPAP is indicated, the patie	ent has a wo	king electr	ical outlet by their bed.	
□ Yes □ No				

TENS UNIT

Criteria for Authorization

May be indicated for patients with chronic pain disorders who are refractory to other treatment and who have demonstrable relief from a TENS trial. No TENS unit will be authorized without a trial.

A completed referral must fully document indications for a TENS unit.

- refer patient to PT for TENS trial with two visits AND
- PT to report results to MMS Staff

THYROID DISEASE

Criteria for Authorization

Hypothyroidism is evaluated and treated at the primary care level. Referral to Endocrinology is approved for severe cases or when the primary care physician is unable to control the disease.

Most patients with hyperthyroidism should be referred to endocrinology to be evaluated and to explore treatment options. Hyperthyroidism secondary to taking excessive thyroid medication is managed by primary care.

Thyroid nodules or other thyroid masses are referred to endocrinology and to either interventional radiology or general surgery for biopsy or other surgery. Occasionally, referral to endocrinology is needed to evaluate and recommend treatment for severely ill or cardiac patients.

Patient History (Indications for Referral)

- for fine needle aspiration of solitary nodules
- for treatment of thyroid cancer
- to confirm the diagnosis and treatment plan for hyperthyroid patients
- for radioactive iodine therapy
- when lab values are ambiguous, especially in sick or elderly patients
- uncontrolled hypothroidism

THYROIDECTOMY

Criteria for Authorization

Patient History (one of two)

- family history of thyroid cancer
- recurrent cystic lesions

AND

Physical (one of three)

- presence of a thyroid nodule or mass
- lymphadenopathy or metastasis
- cystic lesion > 4cm

AND

Diagnostic (one of four)

- fine needle aspiration, positive for cancer
- I₁₃₁, scan-positive
- chronic thyroiditis by microsomal antibodies
- ↑ calcitonin levels

TEMPOROMANDIBULAR JOINT DISORDER-TMJ

Criteria for Authorization

TMJ refers to persistent pain and other symptoms such as clicking in the temporomandibular joint of the jaw. This common problem has many causes: arthritis, dental problems, and stress causing grinding or clenching the teeth (bruxism). TMJ is initially evaluated by primary care to determine the most likely cause. Dental referral should be done before a medical specialist if there is evidence of malocclusion or other dental problems. Other possible referrals include counseling, physical therapy and ENT which are approved based on these criteria:

Patient History (two of four)

- pain or difficulty opening mouth
- iaw lockina
- clicking, popping or crepitus sound
- past history of rheumatoid arthritis or osteoarthritis

AND

Physical Exam (one of three)

- presence of facial asymmetry
- limited movement of the jaw
- tenderness and/or crepitation over TMJ joint on palpation

Dental evaluation should be done on most patients.

Failure of Past Treatment (two of three)

- muscle relaxants
- anti-inflammatory agents
- splint/oral appliance

TONSILLECTOMY AND ADENOIDECTOMY (T & A)

<u>Criteria for Authorization</u>

Chronic persistent infection unresponsive to antibiotics and chronic obstruction are the most common reasons for referral to ENT and removal of these glands.

Patient History (one of three)

- repeated episodes of acute tonsillitis (four or more) in past year with failure of resolution despite antibiotic therapy
- persistent obstruction of breathing and swallowing
- recurrent otitis media with persistence of fluid pressure secondary to enlarged adenoids causing obstruction to the eustachian tubes

Physical Exam (one of three)

- markedly enlarged and chronically infected tonsils
- tonsils causing oral obstruction
- peritonsillar abscess
- adenoid obstruction of the eustachian tubes (by imaging)

Adenoidectomy alone (one of three)

- nasal obstruction resulting in sleep apnea
- chronic otitis media with effusion secondary to adenoids

TRIGGER FINGER

Criteria for Authorization

Trigger finger or stenosing tenosynovitis is a condition in which one or more fingers (including the thumb) is/are caught in a bent position. This finger may straighten with a snap like a trigger being pulled and released. The finger remains in a bent and locked position in more severe cases. Referral and surgery is approved when correction of the trigger finger is critically necessary for work or daily function.

Patient History (one of first two)

- pain at the interphalangeal joint of forefinger or thumb
- failure of injectable steroids
- affecting work (obtain work history) required

AND

Physical Exam (one of two)

- nodular thickening at the M.C.P. joint
- catching or locking of the P.I.P. joint with extension of finger

TYMPANOPLASTY

Criteria for Authorization

Tympanoplasty is repair of the tympanic membrane, or eardrum. The procedure is done for persistent perforations of the eardrum.

Patient History (two of three)

- recurrent infection of the middle ear
- chronic hearing loss interfering with work or daily function
- failure of previous antibiotic therapy and observation to result in healing of the perforation
- trauma resulting in perforation of the eardrum

AND

Physical Exam and Testing (two of two)

- perforation of tympanic membrane
- hearing loss of > 40 db by audiometry

TYMPANOTOMY

Criteria for Authorization

Tympanotomy, also known as myringotomy, is a surgical incision of the tympanic membrane, or eardrum. The procedure is done to perform surgery in the middle ear, or more commonly to insert drainage tubes because of persistent fluid in the middle ear.

Patient History or medical records documenting one of the following:

- cholesteotoma (collection of tissue in the middle ear)
- for insertion of typanostomy tubes (documented need by consult)
- to explore the middle ear for hearing loss (> 40 db) or other pathology

VARICOSE VEINS

Criteria for Authorization

Varicose veins are veins that become enlarged or twisted. Usually these occur in superficial veins of the leg, especially in women during and after pregnancy. Most varicose veins cause no significant medical problems and treatment is not necessary. CMS approves referral and surgery for varicose veins that cause major problems with work or daily function, and never for cosmetic purposes.

Patient History (both present)

- associated with severe, constant pain and/or stasis ulceration
- prescription compression stockings have failed after at least a six-month trial

Note: Patient unlikely to require coronary artery bypass grafting in the future.

Physical Exam (one of these present)

- recurrent superficial phlebitis (two or more occasions)
- stasis ulcer that is recurrent (three or more occasions) or not responding to conservative therapy after six weeks

Contraindication: Occlusive arterial disease (moderate to severe)

- recent deep vein thrombophlebitis
- pregnancy
- congenital abnormalities of deep veins

VERTIGO

Criteria for Authorization

Vertigo is dizziness associated with a feeling of movement, such as the room spinning. Vertigo is usually caused by a problem with the inner ear balance mechanism (vestibular system), or in the brain. The most common cause of vertigo is benign positional vertigo (BPV), a temporary condition common in middle age and the elderly. Temporary vertigo is also caused by inner ear infections, usually a virus, called labyrinthitis. Other more serious causes include toxicity with medications, ischemia to the brain (TIA or stroke) or brain tumors. If the vertigo is caused by an inner ear problem, it is referred to as peripheral vertigo. If the cause is in the brain, it is referred to as central vertigo. The initial assessment of vertigo, including maneuvers to determine if it is peripheral or central, is done by primary care. Referrals for peripheral vertigo usually go to ENT, while central vertigo is referred to Neurology.

Patient History

- True rotatory vertigo elicited by a rapid head movement in a non-axial plane, e.g. rolling over in bed.
- If other neurologic symptoms are present, such as weakness, severe headache or hearing loss, early referral is indicated.
- Failure of treatment for BPV including:
 - medications
 - epley Maneuvers
- Assume position of Dix-Hallpike with the affected ear down then slowly rotate head in the opposite direction. Then, rotate head and whole body another 90 degrees, resume sitting.

AND

Physical Exam

Dix-Hallpike maneuver

VERTIGO (Continued)

- Patient moves from a sitting to a supine position with the head hanging over the edge of the bed or table and rotated 45 degrees; ear down. Bi-lateral testing.
- Affected ear facing ground > vertigo and rotating movement of eyes or nystagmus indicated peripheral vertigo and most likely BPV.
- If this is negative, a central cause of vertigo is considered and neurologic testing is indicated.

Diagnostic Testing

- Audiometry should be done to document hearing loss
- A CT scan (for acoustic neuroma) or MRI (for brain tumor or mass)may be requested by primary care.

San Diego County CMS Program WORK HISTORY INFORMATION

The CMS Program policy limits dental services, specifically stay-plates and dentures. We require specific information from the patient to determine if the requested service meets all of the CMS Program guidelines to be a covered service. Your cooperation is appreciated. **ALL questions must be answered and the form attached to the request for dental replacements.**

Patient Name:	SSN:			
Phone Number:	DOB:			
What kind of dental service do you need?				
When were your teeth extracted? Month		Year		
3. What kind of work do you do when you are working?				
4. Are you currently employed?	[]Yes	[] No		
5. Are you currently Receiving State Disability?	[]Yes	[] No		
6. Are you currently receiving workers compensation?	[]Yes	[] No		
7. Date you last worked?				
IF YOU ARE CURRENTLY UNEMPLOYED:				
Why did you leave your last job?				
2. Have you applied for or been offered employment in the	past (6) months?		[]Yes	[] No
3. Have you recently been turned down for a job because of this medical condition?			[]Yes	[] No
TELL US WHO YOUR CURRENT EMPLOYER IS OR ABO OFFERED YOU EMPLOYMENT	UT THE COMPA	NY WHO HAS		
Name of Company:				
Person to Contact:	Phone:			
If you are currently employed you can speed up the review pemployer and send a letter on <u>business letterhead.</u> This lett employment and how this condition affects your ability to do history and send them to: CMS Program ATTN: Authorization Coord PO Box 939016	orocess if you wo ter should tell us your job. Attach dinators	ould have your about your the letter to thi	s work	
San Diego, CA 9219 I authorize the CMS Program to contact the persons/organiz	zalions nameu al	•		
		•		

WOUND MANAGEMENT

Criteria for Authorization

Patients with diminished circulation or low oxygen in the blood may have chronic wounds that if not treated aggressively become more complicated. Most wound management is done by primary care, but complicated wounds may require the evaluation and management recommendations of a wound care specialist or clinic. CMS authorizes such a referral and treatment procedures if critically necessary for wound healing.

Patient History (all must be present)

- chronic ulcers-not healed within 30 days of occurrence
- failure of standard wound therapy
- no measurable signs of healing

Physical Exam

 chronic stage 3 & 4 pressure ulcers, arterial ulcers, diabetic ulcers and venous stasis